



nutrients



Article

Associations Between Metabolic Age, Sociodemographic Variables, and Lifestyle Factors in Spanish Workers

Ignacio Ramírez-Gallegos, Marta Marina-Arroyo, Ángel Arturo López-González, Daniela Vallejos, Emilio Martínez-Almoyna-Rifá, Pedro Juan Tárraga López and José Ignacio Ramírez-Manent

Special Issue

Effects of Nutrient Intake on Cardiovascular Disease

Edited by
Prof. Dr. Noreen F. Rossi



<https://doi.org/10.3390/nu16234207>

Article

Associations Between Metabolic Age, Sociodemographic Variables, and Lifestyle Factors in Spanish Workers

Ignacio Ramírez-Gallegos ¹, Marta Marina-Arroyo ¹ , Ángel Arturo López-González ^{1,2,3,4,*} , Daniela Vallejos ¹, Emilio Martínez-Almoyna-Rifá ¹, Pedro Juan Tárraga López ^{5,6}  and José Ignacio Ramírez-Manent ^{1,3,4,7} 

- ¹ ADEMA-Health Group University Institute of Health Sciences Research (IUNICS), 07009 Palma, Balearic Islands, Spain; ignacioramirezgallegos@gmail.com (I.R.-G.); marinaarroyomarta@gmail.com (M.M.-A.); d.vallejos@eua.edu.es (D.V.); emilio@udemax.com (E.M.-A.-R.); joseignacio.ramirez@ibsalut.es (J.I.R.-M.)
- ² Faculty of Dentistry, University School ADEMA, 07009 Palma, Balearic Islands, Spain
- ³ Balearic Islands Institute of Health Research (IDISBA), Balearic Islands Health Research Institute Foundation, 07010 Palma, Balearic Islands, Spain
- ⁴ Balearic Islands Health Service, 07010 Palma, Balearic Islands, Spain
- ⁵ Faculty of Medicine, University of Castilla la Mancha, 02008 Albacete, Castilla-La Mancha, Spain; pjtarraga@sescam.jccm.es
- ⁶ SESCOAM (Servicio Salud Castilla La Mancha), 45071 Toledo, Castilla-La Mancha, Spain
- ⁷ Faculty of Medicine, University of the Balearic Islands, 07010 Palma, Balearic Islands, Spain
- * Correspondence: angarturo@gmail.com

Abstract: Background: Metabolic age is defined as an estimation of a person's age based on their basal metabolic rate (BMR) and other physiological health indicators. Unlike chronological age, which simply measures the number of years lived since birth, metabolic age is based on various health and fitness markers that estimate the body's "true" biological age and can be assessed using various methodologies, including bioimpedance. The aim of this study was to evaluate how age, sex, social class, smoking habits, physical activity, and adherence to the Mediterranean diet influence metabolic age. Methods: A cross-sectional, descriptive study was conducted on 8590 Spanish workers in the Balearic Islands. A series of sociodemographic variables and health-related habits were assessed, while metabolic age was measured using bioimpedance. A metabolic age exceeding chronological age by 12 years or more was considered high. A descriptive analysis of categorical variables was performed by calculating their frequency and distribution. By applying multivariate models, specifically multinomial logistic regression, we observe that all independent variables (sex, age, social class, physical activity, mediterranean diet, and smoking) show varying levels of association with the occurrence of high metabolic age values. Among these independent variables, those showing the highest degree of association, represented by odds ratios, are physical activity, adherence to the Mediterranean diet, and social class. In all cases, the observed differences demonstrate a high level of statistical significance ($p < 0.001$). Results: The factors with the greatest influence were physical inactivity, with an OR of 5.07; and low adherence to the Mediterranean diet, with an OR of 2.8; followed by social class, with an OR of 2.51. Metabolic age increased with chronological age and was higher in males, with an OR of 1.38. Smoking also had a negative impact on metabolic age, with an OR of 1.19. Conclusions: Mediterranean diet is associated with a higher metabolic age. The most influential factors on metabolic age are physical activity and adherence to the Mediterranean diet, followed by the individual's socioeconomic class. Smoking also contributes to increased metabolic age, albeit to a lesser extent.



Citation: Ramírez-Gallegos, I.; Marina-Arroyo, M.; López-González, Á.A.; Vallejos, D.; Martínez-Almoyna-Rifá, E.; Tárraga López, P.J.; Ramírez-Manent, J.I. Associations Between Metabolic Age, Sociodemographic Variables, and Lifestyle Factors in Spanish Workers. *Nutrients* **2024**, *16*, 4207. <https://doi.org/10.3390/nu16234207>

Academic Editor: Noreen F. Rossi

Received: 19 November 2024

Revised: 1 December 2024

Accepted: 4 December 2024

Published: 5 December 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: metabolic age; mediterranean diet; physical activity; smoking; social class; lifestyle

1. Introduction

Metabolic age is a relatively new concept in the field of health and wellness. It is defined as an estimate of a person's age based on their basal metabolic rate (BMR) and

other indicators of physiological health, providing an assessment of the body's 'true' biological age [1,2].

Metabolic age is a key indicator of physical fitness, and a lower metabolic age is associated with better physical fitness and a longer, healthier life. Metabolic age is influenced by factors such as genetics, body composition, physical activity, and diet. It alters BMR, which is affected by muscle mass [3], genetic factors [4,5], sex [6], age [7,8], chronic diseases [9], diet [10–12], stress [13,14], and sleep quality [15,16]. It has significant implications for both individual health and healthcare costs. When a person's metabolic age exceeds their chronological age, potential years of health may be lost. The concept of Avoidable Lost Life Years (ALLY) is introduced as a useful measure to assess this loss.

Interventions to improve metabolic age include lifestyle changes, such as increased physical activity, a balanced diet, and stress management [17,18]. Resistance training can be particularly effective for increasing muscle mass and improving BMR [19]. Additionally, nutritional interventions that promote body fat loss and metabolic health improvement are also recommended [20].

Metabolic age can be assessed using various methodologies. One of the most common is the use of bioimpedance scales which estimate body composition and BMR [21]. These scales send a low-intensity electrical current through the body to measure the resistance of different tissues [22].

Another tool used is indirect calorimetry, which measures oxygen consumption and carbon dioxide production to calculate BMR [23,24]. Blood tests evaluating metabolic markers, such as glucose and lipid levels, can also provide valuable information about metabolic health [25].

Having a metabolic age significantly higher than the chronological age can be an indicator of a higher risk of chronic diseases, such as type 2 diabetes, cardiovascular diseases, and certain types of cancer [26]. On the other hand, a metabolic age lower than the chronological age can be a sign of good health and a lower risk of these diseases [27].

Identifying the variables that most influence the metabolic age of our population can be highly valuable for implementing health interventions aimed at reducing metabolic age, improving quality of life within this population, and lowering healthcare costs.

The objective of this study was to evaluate how different variables, both sociodemographic (age, sex, and social class) and health-related habits (smoking, physical activity, and Mediterranean diet), affect metabolic age values determined by bioimpedance in a group of workers in the Balearic Islands.

2. Materials and Methods

2.1. Participants

A cross-sectional, descriptive study was conducted on 8590 Spanish workers in the Balearic Islands. The sample consisted of workers who attended their annual occupational health check-up between January 2019 and December 2020 at an occupational health and risk prevention service. This service caters to a range of companies, notably in the healthcare, public administration, hospitality, retail, transportation, education, industrial, and cleaning sectors.

Refer to the flow chart in Figure 1 for details.

Inclusion criteria:

- Individuals aged 18 to 69 years.
- Willingness to participate in the research.
- Consent for the data to be used in epidemiological studies.
- Employment with one of the companies participating in the research and not being temporarily disabled at the time of the study.

Exclusion criteria:

- Age under 18 years or over 69 years.
- Not being an employee of one of the participating companies.

- Refusal to participate in the research study.
- Refusal to consent to the use of data for epidemiological studies.
- Lacked a parameter for calculating scales.




Figure 1. PRISMA flowchart of participants in this study.

2.2. Determination of Variables

The health personnel from the occupational health services of the participating companies were responsible for gathering the necessary data for this study through:

Anamnesis: A detailed clinical history was compiled, which included information on sociodemographic variables such as age, sex, social class, tobacco consumption, physical exercise, and adherence to the Mediterranean diet.

Anthropometric and clinical measurements: These included height, weight, waist and hip circumference, and systolic and diastolic blood pressure.

Analytical determinations: The lipid profile and blood glucose levels were measured.

2.2.1. Anthropometric Determinations

To minimize potential biases in the study, measurement techniques for the variables were standardized. Height and weight were measured using a SECA 700 scale and a SECA 220 stadiometer. Measurements were taken with the individual in underwear, following international standards for anthropometric evaluation as outlined by ISAK [28]. Data were recorded in centimeters and kilograms.

For the assessment of abdominal waist circumference, a SECA measuring tape was used, placed midway between the last rib and the iliac crest, parallel to the floor. The individual stood with their abdomen relaxed. Hip circumference was measured similarly, with the tape measure parallel to the floor at the widest part of the buttocks [29].

2.2.2. Clinical Determinations

Blood pressure was measured with an OMRON-M3 model blood pressure monitor. For accurate assessment, the individual sat with their back supported by the chair and rested for at least 10 min. The patient needed to be relaxed, with their arm supported at heart level, without crossing their legs. They should not have eaten, smoked, or consumed alcohol or tea for at least an hour before the measurement. The cuff was placed around the upper arm, 2–3 cm above the elbow crease, ensuring a snug fit without being too

tight. Various cuff sizes were available. Three consecutive measurements were taken at one-minute intervals, with the final value being the average of the three.

2.2.3. Analytical Determinations

Blood samples were collected via venipuncture after a 12 h fast. Samples were then processed and refrigerated to ensure proper preservation for no more than 48–72 h. Analysis of the samples was conducted in reference laboratories using standardized methodologies. Triglycerides, total cholesterol, and blood glucose were measured using enzymatic methods, while HDL cholesterol was assessed using precipitation methods. LDL cholesterol was calculated indirectly using the Friedewald formula, valid when triglyceride levels did not exceed 400 mg/dL. If triglyceride levels were greater than 400 mg/dL, LDL was measured directly. All analytical variables are expressed in mg/dL.

2.2.4. Risk Scales

Adherence to the Mediterranean diet was evaluated using the PREDIMED questionnaire [30], which consists of 14 questions, each scoring 0 or 1. A score of nine or more indicates high adherence to the diet [31].

The level of physical activity was measured using the International Physical Activity Questionnaire (IPAQ), a self-administered survey that evaluates physical activity over the previous seven days [32].

A person who had smoked at least one cigarette per day (or its equivalent in other forms of consumption) in the past thirty days, or who had quit smoking less than twelve months before, was considered a smoker. A person who had not smoked in the past year or who had never smoked was considered a non-smoker.

Determination of socioeconomic class followed the recommendation of the Spanish Society of Epidemiology, based on the 2011 National Classification of Occupations [33]. According to this classification: Class I includes managers, directors, and university professionals; Class II comprises intermediate professions and self-employed individuals; and Class III consists of manual laborers.

Metabolic age was measured using a TANITA MC-780 S MA bioimpedance meter (TANITA Corporation, Tokyo, Japan).

ALLY (Avoidable Lost Life Years) was calculated by subtracting metabolic age from biological age. Some published studies have reported that a difference of at least 12 years between chronological and metabolic age reduces cardiovascular risk. ALLY is classified as low if the difference is less than 3 years, normal if it is between 3 and 11 years, and high if the difference is 12 years or more. A metabolic age 12 years or more above the chronological age was considered high [34]. This serves as the cutoff point for establishing high metabolic age values for ALLY.

2.3. Statistical Analysis

A descriptive analysis of categorical variables was performed, calculating their frequency and distribution. For quantitative variables with a normal distribution, the mean and standard deviation were calculated. To compare means, Student's *t*-test was used, and to compare proportions, the chi-square test was used. The dependent variable, ALLY, was classified into three categories. Since it comprises more than two categories, we performed a multivariate analysis stratified by age groups to assess the influence of each studied variable within each stratum. The independent variables selected were those considered, according to the reviewed literature, to be the most statistically and biologically recommended. The multivariate analysis was conducted using multinomial logistic regression, along with the calculation of odds ratios and the Hosmer–Lemeshow goodness-of-fit test. SPSS 29.0 software was used for the statistical analysis, with an accepted level of statistical significance of 0.05.

3. Results

The anthropometric and clinical details of the study participants are presented in Table 1. The analysis included a total of 8590 individuals (4104 men, 47.8%, and 4486 women, 52.2%). The average age of the sample was slightly over 41 years, with the majority of participants between 30 and 49 years old. The analysis of anthropometric, clinical, and analytical variables revealed significantly lower values in females across all measures. The majority of the sample subjects belonged to social class I. In both sexes, just over 15% were smokers. Among men and women, 25.9% and 35.1%, respectively, did not engage in regular physical exercise. Additionally, more than half of the sample in both sexes adhered to the Mediterranean diet.

Table 1. Characteristics of the participants.

	Men n = 4104	Women n = 4486	
	Mean (SD)	Mean (SD)	p-Value
Age (years)	41.6 (10.6)	41.5 (10.5)	0.492
Height (cm)	175.8 (7.2)	162.5 (6.1)	<0.001
Weight (kg)	81.2 (14.8)	63.9 (13.6)	<0.001
Waist circumference (cm)	89.8 (12.5)	77.0 (12.0)	<0.001
Hip circumference (cm)	101.8 (8.7)	99.6 (10.9)	<0.001
Systolic blood pressure (mmHg)	128.6 (13.3)	117.2 (14.1)	<0.001
Diastolic blood pressure (mmHg)	79.9 (10.2)	74.9 (9.9)	<0.001
Glycaemia (mg/dL)	93.4 (17.8)	88.9 (12.6)	<0.001
Total cholesterol (mg/dL)	191.8 (36.0)	189.0 (34.8)	<0.001
HDL-cholesterol (mg/dL)	49.2 (11.3)	59.5 (12.8)	<0.001
LDL-cholesterol (mg/dL)	124.0 (54.6)	113.8 (30.7)	<0.001
Triglycerides (mg/dL)	107.8 (69.4)	81.5 (46.3)	<0.001
GGT (UI)	31.5 (30.0)	18.5 (15.9)	<0.001
AST (UI)	24.4 (17.3)	18.2 (7.7)	<0.001
ALT (UI)	29.3 (34.9)	17.3 (13.4)	<0.001
	%	%	p-value
18–29 years	15.5	16.8	0.005
30–39 years	27.8	25.1	
40–49 years	32.7	34.4	
50–59 years	19.0	19.7	
60–69 years	5.0	4.0	
Social class I	57.1	50.8	<0.001
Social class II	20.2	23.8	
Social class III	22.7	25.4	
Non-smokers	84.5	84.2	0.348
Smokers	15.5	15.8	
Physical inactivity	25.9	35.1	<0.001
PhA Moderate	27.0	26.5	
PhA High	47.1	38.4	
NAD Mediterranean diet	44.5	41.6	<0.001
AD Mediterranean diet	55.5	58.4	

SD Standard deviation. HDL-c High-density lipoprotein. LDL-c Low-density lipoprotein. GGT Gamma-Glutamyl Transferase. AST Aspartate Aminotransferase. ALT Alanine Aminotransferase. PhA Physical activity. AD Mediterranean diet: Adherence to the Mediterranean diet. NAD Mediterranean diet: No Adherence to the Mediterranean diet. Student’s *t*-test was used for means and chi square test for prevalence.

Table 2 presents the mean values of the ALLY metabolic age, stratified by age and various variables for each sex. Our results include several negative values, indicating that the metabolic age is lower than chronological age, particularly noticeable in the youngest age group, 18 to 29 years. In nearly all values obtained, the mean metabolic age is lower in women compared to men. In both sexes, the ALLY metabolic age averages worsen with decreasing socioeconomic status, smoking, lack of regular physical activity, and low adherence to the Mediterranean diet. Notably, regular physical activity, practiced at least

three days a week, emerges as the variable most significantly associated with a lower metabolic age across all ages and in both sexes. In all the results, the observed differences are statistically significant ($p < 0.001$).

Table 2. Mean metabolic age values stratified by age and according to different sociodemographic variables and healthy habits by sex.

		18–29 Years		30–39 Years		40–49 Years		50–59 Years		60–69 Years	
Men	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	
Social class I	474	−5.2 (10.5) *	774	−6.6 (9.6) *	630	−5.5 (10.6) *	324	−3.8 (10.8) *	144	−3.9 (11.1) *	
Social class II	84	−3.7 (6.7) *	204	−0.1 (10.9) *	300	−3.5 (11.3) *	216	−3.4 (9.9) *	24	2.0 (11.0) *	
Social class III	78	−2.5 (10.4) *	162	1.4 (13.3) *	414	−2.4 (11.2) *	240	0.7 (12.5) *	36	8.0 (7.2) *	
Non-smokers	534	−5.7 (9.6) *	924	−4.4 (11.1) *	1158	−4.6 (10.9) *	672	−2.5 (11.1) *	180	−1.5 (11.5) *	
Smokers	102	0.5 (11.2) *	216	0.2 (10.6) *	186	−0.7 (11.3) *	108	1.3 (12.2) *	24	1.3 (11.1) *	
Physical inactivity	120	1.7 (11.2) *	270	1.4 (11.9) *	390	3.7 (10.8) *	198	3.8 (9.9) *	84	6.0 (9.6) *	
PhA Moderate	144	−6.3 (8.8) *	348	−2.8 (10.7) *	294	−4.6 (8.5) *	264	−0.7 (11.2) *	60	−5.5 (8.7) *	
PhA High	372	−8.2 (9.5) *	522	−8.3 (8.9) *	660	−8.5 (9.6) *	318	−7.4 (9.8) *	60	−7.9 (10.2) *	
NAD Mediterranean diet	225	−1.4 (11.3) *	426	−1.1 (12.1) *	610	−0.2 (12.3) *	433	1.4 (11.4) *	133	3.4 (10.6) *	
AD Mediterranean diet	411	−6.4 (8.9) *	714	−6.2 (9.8) *	734	−7.3 (8.6) *	347	−6.9 (9.3) *	71	−10.6 (6.1) *	
Women	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	
Social class I	606	−6.2 (9.2) *	702	−8.0 (9.7) *	682	−8.0 (9.5) *	222	−7.9 (10.2) *	66	−7.0 (10.5) *	
Social class II	80	−3.6 (10.5) *	192	−3.9 (10.5) *	464	−2.7 (12.3) *	304	−1.9 (11.9) *	28	−1.2 (12.7) *	
Social class III	68	−2.1 (10.8) *	232	−1.7 (11.9) *	398	−1.5 (12.2) *	356	0.6 (11.5) *	86	0.9 (11.3) *	
Non-smokers	694	−5.3 (9.8) *	942	−6.2 (10.5) *	1304	−5.0 (11.3) *	682	−6.5 (10.2) *	154	−5.0 (11.9) *	
Smokers	60	−4.3 (10.5) *	184	−3.6 (11.3) *	240	−3.3 (12.5) *	200	−2.0 (10.4) *	26	0.2 (10.7) *	
Physical inactivity	192	−3.1 (11.3) *	366	−1.1 (11.5) *	600	0.3 (12.1) *	356	0.5 (12.2) *	60	0.6 (12.1) *	
PhA Moderate	238	−5.5 (8.8) *	330	−6.9 (9.7) *	376	−5.6 (10.5) *	209	−5.1 (10.9) *	34	−7.2 (11.0) *	
PhA High	324	−6.2 (9.5) *	430	−9.5 (9.0) *	568	−9.5 (9.0) *	317	−8.6 (9.5) *	86	−10.4 (8.3) *	
NAD Mediterranean diet	297	−3.2 (11.0) *	425	−4.9 (11.3) *	653	−2.7 (12.1) *	399	−2.5 (12.4) *	92	−0.9 (12.0) *	
AD Mediterranean diet	457	−6.5 (8.8) *	701	−6.6 (10.2) *	891	−6.2 (10.7) *	483	−6.7 (10.2) *	88	−8.7 (10.0) *	

PhA Physical activity. AD Mediterranean diet: Adherence to the Mediterranean diet. NAD Mediterranean diet: No Adherence to the Mediterranean diet. SD Standard deviation. (*) $p < 0.001$. Student’s *t*-test was used.

Table 3 presents the prevalence of high ALLY metabolic age values. When comparing between sexes, the results consistently show a lower percentage in women, suggesting that the studied variables may have less influence on the ALLY metabolic age in females. Comparing across age groups reveals an increase in the percentage of individuals with high metabolic age values with advancing age, reflecting trends related to socioeconomic status and unhealthy lifestyle habits. This pattern is consistent across both sexes. Interestingly, a similar trend is observed for variables associated with healthy lifestyle behaviors (e.g., non-smoking, engaging in regular physical activity three or more days per week, and adherence to the Mediterranean diet). This may indicate that the influence of these factors on metabolic age diminishes with age or that their effects are confounded by interactions between variables. In all analyzed outcomes, the differences observed were statistically significant ($p < 0.001$).

We performed a multinomial logistic regression analysis stratified by age groups to assess the effect of sex, social class, physical activity, adherence to the Mediterranean diet, and tobacco consumption on the likelihood of presenting with a high ALLY metabolic age (dependent variable), as shown in Table 4. This approach allowed us to evaluate the risk associated with each variable independently. Across all age groups, we observed gender differences, with a higher risk of high ALLY metabolic age in men compared to women. Although all modifiable variables showed an increased risk for high ALLY metabolic age, the most influential factors (highest odds ratios) were physical activity and adherence to the Mediterranean diet. Therefore, it is crucial to focus efforts on improving adherence to these two factors in primary care consultations, without neglecting the others. In all the factors studied, the observed differences were statistically significant ($p < 0.001$).

Table 3. Prevalence of high metabolic age values stratified by age and according to different sociodemographic variables and healthy habits by sex.

	18–29 Years		30–39 Years		40–49 Years		50–59 Years		60–69 Years	
Men	n	%	n	%	n	%	n	%	n	%
Social class I	474	13.2 *	774	15.5 *	630	20.8 *	324	22.2 *	144	24.8 *
Social class II	84	21.5 *	204	25.3 *	300	28.0 *	216	30.9 *	24	34.5 *
Social class III	78	23.1 *	162	28.1 *	414	34.8 *	240	35.4 *	36	39.9 *
Non-smokers	534	20.7 *	924	23.5 *	1158	25.9 *	672	27.7 *	180	30.9 *
Smokers	102	28.9 *	216	28.8 *	186	32.7 *	108	33.3 *	24	38.9 *
Physical inactivity	120	40.0 *	270	44.4 *	390	46.9 *	198	50.3 *	84	54.6 *
PhA Moderate	144	14.5 *	348	17.6 *	294	18.4 *	264	21.8 *	60	26.8 *
PhA High	372	12.2 *	522	12.6 *	660	14.5 *	318	15.8 *	60	18.2 *
NAD Mediterranean diet	225	32.0 *	426	37.1 *	610	40.8 *	433	42.3 *	133	44.9 *
AD Mediterranean diet	411	15.7 *	714	19.4 *	734	21.5 *	347	24.2 *	71	27.9 *
Women	n	%	n	%	n	%	n	%	n	%
Social class I	606	10.5 *	702	12.2 *	682	13.8 *	222	16.2 *	66	18.2 *
Social class II	80	19.2 *	192	20.8 *	464	24.7 *	304	27.6 *	28	28.6 *
Social class III	68	20.6 *	232	26.8 *	398	28.9 *	356	29.2 *	86	31.8 *
Non-smokers	694	19.3 *	942	18.9 *	1304	20.5 *	682	23.3 *	154	23.4 *
Smokers	60	23.3 *	184	23.9 *	240	27.8 *	200	29.9 *	26	32.3 *
Physical inactivity	192	29.2 *	366	34.4 *	600	38.8 *	356	43.0 *	60	43.3 *
PhA Moderate	238	12.8 *	330	14.9 *	376	19.1 *	209	21.1 *	34	23.5 *
PhA High	324	8.7 *	430	10.7 *	568	11.6 *	317	13.2 *	86	14.6 *
NAD Mediterranean diet	297	20.3 *	425	22.6 *	653	30.5 *	399	33.3 *	92	38.3 *
AD Mediterranean diet	457	14.6 *	701	18.0 *	891	19.9 *	483	21.8 *	88	23.2 *

PhA Physical activity. AD Mediterranean diet: Adherence to the Mediterranean diet. NAD Mediterranean diet: No Adherence to the Mediterranean diet. (*) $p < 0.001$. The chi-square test was used.

Table 4. Multinomial logistic regression stratified by age.

	18–29 Years	30–39 Years	40–49 Years	50–59 Years	60–69 Years
ALLY Metabolic Age High	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Women	1	1	1	1	1
Men	1.33 (1.28–1.39) *	1.74 (1.50–1.98) *	1.36 (1.29–1.44) *	1.29 (1.23–1.35) *	1.24 (1.20–1.29) *
Social class I	1	1	1	1	1
Social class II	1.19 (1.16–1.23) *	1.59 (1.45–1.74) *	1.71 (1.54–1.89) *	2.47 (1.99–2.96) *	1.45 (1.31–1.59) *
Social class III	1.54 (1.40–1.68) *	3.54 (2.71–4.34) *	2.58 (2.06–3.11) *	2.74 (2.03–3.45) *	1.56 (1.38–1.75) *
Non-smokers	1	1	1	1	1
Smokers	2.13 (1.82–2.45) *	1.45 (1.38–1.53) *	1.14 (1.10–1.17) *	1.31 (1.26–1.37) *	1.23 (1.19–1.27) *
PhA High	1	1	1	1	1
PhA Moderate	2.69 (2.28–3.09) *	2.12 (1.78–2.46) *	3.39 (2.65–4.14) *	1.61 (1.39–1.83) *	3.86 (2.99–4.74) *
Physical inactivity	2.99 (2.14–3.85) *	4.60 (3.51–5.70) *	7.11 (5.67–8.55) *	4.54 (3.36–5.73) *	8.36 (6.90–9.82) *
AD Mediterranean diet	1	1	1	1	1
NAD Mediterranean diet	2.84 (2.13–3.55) *	1.84 (1.48–2.20) *	3.33 (2.75–3.89) *	3.39 (2.63–4.15) *	4.44 (3.05–5.84) *

PhA Physical activity. AD Mediterranean diet: Adherence to the Mediterranean diet. NAD Mediterranean diet: No Adherence to the Mediterranean diet. OR: Odds ratio. CI: Confidence interval. Statistical differences (*) $p < 0.001$.

4. Discussion

Biological aging and chronological age are closely related but not synonymous. Advanced age is associated with decreased functional capacity, a chronic hyperinflammatory state, and a higher prevalence of chronic diseases, which may accelerate metabolic age [35]. Previous studies have shown that advanced chronological age is associated with a higher risk of metabolic diseases and increasing metabolic age [36]. Moreover, deterioration of the endocrine and immune systems, along with decreased muscle mass and increased fat mass, are characteristics of aging that contribute to a higher metabolic age. However, biological aging is not a disease but rather an ascending process that occurs progressively in all living beings that surpass the so-called essential life span (ELS) of their species. For humans, this ELS point is typically between 40 and 50 years of age. An individual’s ability to survive

beyond this point serves as an indicator of their health status, their capacity to endure, and ability to maintain a healthy functional state. Therefore, it is crucial to identify the factors that influence an individual's metabolic age in order to intervene in their homeodynamics and enhance their functional status and quality of life [37].

In our sample, metabolic age increased with chronological age, lower socioeconomic status, smoking, sedentary behavior, and low adherence to the Mediterranean diet. Women consistently showed lower metabolic age compared to men across all variables. Among the studied factors, age had the most significant impact on metabolic age, reflecting the natural decline in biological maintenance and repair systems as part of the aging process [38].

Sex also plays a crucial role in determining metabolic age. Men generally have greater muscle mass and lower fat mass compared to women, which could initially be advantageous in terms of basal metabolism. However, men are also more predisposed to accumulating visceral fat, which is associated with a higher risk of metabolic and cardiovascular diseases [39]. Women, especially after menopause, undergo hormonal changes, such as reduced estrogen levels, which can lead to increased abdominal fat and decreased muscle mass, contributing to a higher metabolic age [40]. In our study, women exhibited a lower risk of high ALLY metabolic age compared to men across all age groups. In our study, we did not account for the phase of the menstrual cycle, during which Kanellakis et al. [41] observed changes in body composition and approximately 0.5 kg of weight variation, primarily due to extracellular fluid retention during menstruation. We acknowledged this limitation, as more than half of our sample consisted of women of reproductive age.

In our study, socioeconomic level emerged as another significant determinant of metabolic health. Our findings indicate that ALLY (Avoidable Lost Life Years) increases as socioeconomic class decreases. Specifically, in social class I, values across all age groups are negative, indicating a metabolic age lower than chronological age. In contrast, in social class III, values are less negative, and in three age groups, they even become positive, with the largest differences observed in the 60–69 age group for both men and women. We observed similar trends in the prevalence of high metabolic age values, with percentages increasing alongside the individuals' age. Multinomial logistic regression reveals an elevated risk of high ALLY metabolic age according to socioeconomic class. Therefore, in our results, socioeconomic class also shows a strong association with metabolic age. Although we did not find other studies establishing this association, we believe it could be influenced by the fact that individuals in lower socioeconomic classes often have limited access to health resources, education, and healthy food, potentially impacting their metabolic age negatively [42]. Studies have shown that stress associated with disadvantaged socioeconomic conditions can lead to unhealthy behaviors [43], such as poor diet and lack of physical activity, which increase the risk of metabolic diseases. Additionally, limited access to preventive health services and adequate treatments can exacerbate these conditions [44].

Smoking is a well-known risk factor for numerous chronic diseases, including metabolic and cardiovascular conditions [45]. Nicotine and other toxic substances in tobacco negatively affect the metabolism by increasing insulin resistance, causing systemic inflammation, and altering lipid profiles [46]. Smoking is also associated with reduced muscle mass and increased visceral fat, factors that contribute to a higher metabolic age [47]. Various studies have shown that smokers have a higher risk of developing type 2 diabetes and metabolic syndrome, which accelerates their metabolic aging [48]. Our study found that smoking significantly impacted metabolic age, with a substantial ALLY index that was more pronounced in men but statistically significant for both sexes. High metabolic age was more prevalent among smokers of both genders, with metabolic age progressively increasing with age in the smoking group. This trend was consistently higher in men across all age groups. Multinomial logistic regression analysis confirmed that smoking significantly increased the risk of high metabolic age across all age groups.

Regular physical activity has been shown to reduce the risk of at least 26 chronic diseases, promoting health and quality of life. Similarly, a lack of physical activity has been

linked to an increased risk of disease. Studies demonstrate that the more regular physical activity or exercise one engages in, the greater the health benefits. Moreover, regular physical activity can induce metabolic changes that lead to improved cardiometabolic health [49]. Thus, regular physical activity is one of the most effective factors in reducing metabolic age. Exercise improves insulin sensitivity, reduces visceral fat, and increases muscle mass, all of which are beneficial to metabolic health [50]. Additionally, regular physical activity reduces systemic inflammation and improves lipid profiles, which can counteract the negative effects of chronological aging [51]. In our study, physical activity was the variable most strongly associated with ALLY. Both in terms of average metabolic age values and the prevalence of high metabolic age, it was the factor that showed the greatest differences between those who did not engage in regular physical activity and those who maintained a high level of exercise. These results were consistent across all age groups and in both men and women. The largest differences were observed in the older age groups for both sexes, which may be related to the importance of lifelong physical exercise in mitigating the effects of biological aging.

Although no single diet is universally beneficial for health, a healthy diet emphasizes following dietary guidelines rather than restricting specific nutrients [52]. In our study, we assessed the Mediterranean diet. This diet, rich in fruits, vegetables, whole grains, olive oil, fish, and nuts, has been associated with numerous benefits for metabolic health. It is characterized by high levels of antioxidants, fiber, and healthy fats, which help reduce inflammation and improve insulin sensitivity [53].

The Mediterranean diet has demonstrated benefits for cardiovascular disease, diabetes mellitus, metabolic syndrome, overweight and obesity, various cancers, cognitive function, and overall mortality reduction. Further, the Mediterranean diet is linked to improved body composition and reduced visceral fat accumulation, key factors for achieving optimal metabolic health [54]. However, to our knowledge, this is the first study linking the Mediterranean diet with metabolic age. Our results show that adherence to a Mediterranean diet has a substantial influence on metabolic age, with the highest OR observed across all age ranges following physical activity.

In the results of our study, we observed that ALLY, in both men and women, showed a significant difference between those who adhered to the Mediterranean diet and those who did not. However, in this case, the differences were greater in men across all age groups compared to women. In the multinomial logistic regression, we found that lack of adherence to the Mediterranean diet increased the risk of high ALLY metabolic age from 184% in the 30–39 age group to 444% in the oldest age group. This makes the Mediterranean diet the second most influential variable in determining an individual's metabolic age.

Given the importance of these findings, it is essential to promote adherence to the Mediterranean diet across all age groups, which requires appropriate health and nutrition promotion policies and interventions. This is particularly urgent, as a systematic review conducted by Obeid et al. in 2022 reported a significant decline in adherence to the Mediterranean diet in Mediterranean Basin countries [55].

Similarly, a study conducted in Spain by Herrera-Ramos et al. in 2023 also identified a decline in adherence to the Mediterranean diet among Spanish children and adolescents over the past 20 years. Although the adherence levels reported in this study were slightly higher than those observed in our study, the decreasing consumption of key Mediterranean foods is concerning. For instance, the frequency of fish consumption two to three times per week dropped from approximately 80% in the 1998–2000 cohort to 65% in the 2019–2020 cohort. Likewise, legume consumption decreased from 90% in the 1998–2000 cohort to about 55% in the 2019–2020 cohort [56].

This study highlights the deterioration of dietary habits among Spanish children and adolescents and underscores the urgent need to ensure a high-quality diet. It is crucial for children and adolescents to adopt a Mediterranean diet, as it benefits their health and quality of life. Furthermore, developing healthy dietary habits during childhood increases the likelihood of their continuation into adulthood.

The global population aged over 65 is increasing, and it is projected to reach 2.1 billion people by 2050 [57]. Aging is associated with increased morbidity and changes in body composition, including an increase in fat mass, a decrease in muscle mass, and a reduction in basal metabolic rate (BMR) [58]. This situation is linked to obesity and numerous related pathologies [59], leading to a significant decline in quality of life and rising costs for society and the healthcare system, which may become unsustainable. Consequently, it is crucial to seek effective strategies that promote healthy aging and help mitigate morbidity, thereby enhancing quality of life during these additional years.

Among these strategies is the potential reduction in metabolic age by targeting various associated factors, which we have evaluated in this study. In this context, it is important to emphasize that a metabolic age lower than chronological age increases BMR, thereby reducing mortality and morbidity [60]. In our study, we examined modifiable variables associated with metabolic age that could be targeted to improve it. These variables were assessed across different age and sex strata to understand how each factor influences these groups and to determine if focusing on specific factors by sex and age group would result in more efficient health interventions.

Public spending increases with age due to the growing health and social care needs of the elderly, which are influenced by their diminished ability to manage daily living activities and reduced physical capacities [61]. Therefore, resource needs for older individuals are more closely related to disability than chronological age. One of the primary determinants of healthcare costs is the health status of the population. Addressing modifiable risk factors associated with non-communicable diseases is a cost-effective strategy to reduce their burden. These risk factors include harmful alcohol consumption, smoking, unhealthy diets, and physical inactivity—all of which also contribute to metabolic age. Every dollar invested in interventions aimed at reducing these risk factors could potentially generate a savings of 7 dollars (USD) [62].

In our study, physical inactivity emerged as the most influential variable affecting metabolic age. A study published in *The Lancet* in 2016 estimated the global health-related costs of physical inactivity at 53.8 billion international dollars (INT), a value representing the amount of goods and services that an individual or government could purchase in their respective country compared to what could be acquired in the United States [63,64]. Another study published in *The Lancet* in 2023 estimated the global cost of physical inactivity to be approximately 520 billion INT\$ over the 2020–2030 decade [65].

Preventing disease earlier in life leads to a better quality of life and lower healthcare costs. Interventions aimed at reducing needs in old age should be implemented across all age groups, as they can benefit the healthcare system. The dissemination and implementation of strategies to improve health are essential to reduce social and healthcare costs. In an era of increasing budgetary constraints, analyzing costs and identifying the factors that influence them could help define strategies for cost reduction.

5. Strengths and Limitations

One of the strengths of the study is the large sample size, nearly 8600 people, and the wide variety of variables analyzed.

The main limitation is that only individuals of working age (18–69 years) were included, thus excluding unemployed individuals, retirees, those under 18, and those over 69 years old. Our results cannot be extrapolated to the entire population as some age groups are missing.

Since more than half of the sample belongs to social class I, it may not be representative of the general population.

Likewise, since this is a population of the Balearic Islands only, it is possible that the results may differ in other types of populations. Therefore, our results cannot be extrapolated to them.

Other confounding factors such as comorbidities or pharmacological treatments were not included, as these data were not available.

Another limitation of our study is the healthy worker bias, which is a common methodological problem in research studies on workers. Workers with chronic diseases or who are more prone to illness may be less likely to attend occupational health check-ups compared to healthy workers, which could underestimate the results.

Since this is a cross-sectional study, it allows for an association between the variables studied and obesity, but we cannot establish causality.

Finally, we have not taken into account the menstrual phase of the women included in the study, which may produce variations in their weight.

6. Conclusions

Metabolic age is shaped by both demographic and lifestyle factors. While chronological age and sex are unmodifiable influences, modifiable factors like physical activity and adherence to the Mediterranean diet have the most significant impact, followed by socioeconomic status and smoking habits. Addressing these modifiable factors through targeted interventions and public health policies is crucial to improving metabolic health, reducing chronic disease burdens, lowering healthcare costs, and enhancing overall quality of life.

Author Contributions: Conceptualization: Á.A.L.-G., J.I.R.-M. and I.R.-G.; Data collection and analysis: Á.A.L.-G., I.R.-G. and E.M.-A.-R.; Data curation: Á.A.L.-G. and M.M.-A.; Methodology: I.R.-G. and Á.A.L.-G.; Validation: M.M.-A., D.V. and P.J.T.L.; Formal analysis: Á.A.L.-G.; Investigation: I.R.-G., D.V. and P.J.T.L.; Draft: I.R.-G. Revision: J.I.R.-M. and Á.A.L.-G. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The research team adhered strictly to national and international ethical guidelines for health sciences research, as established by the Declaration of Helsinki, with special attention given to participant anonymity and data confidentiality. Approval was obtained from the Ethics and Research Committee of the Balearic Islands (CEI-IB) using the indicator IB 4383/20, approval date: 26 November 2020. Participation in the study was voluntary; participants were fully informed about the study's purpose and provided both written and verbal consent. They received an information sheet detailing the study's objectives and an informed consent form. To ensure confidentiality, the data collected was identified using codes that only the project coordinator could link to the participants. Participant identities will not be published in any study report, and no identifying information will be shared by the investigators. The research team guarantees that participants can exercise their rights to access, rectify, cancel, and oppose the data collected. Furthermore, the team undertakes to comply strictly with Organic Law 3/2018, of December 5, on the protection of personal data and guarantee of digital rights.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: This study data are stored in a database that complies with all security measures at ADEMA-Escuela Universitaria. The Data Protection Delegate is Ángel Arturo López González.

Conflicts of Interest: The authors declare that they have no conflicts of interest.

References

1. Johnson, A.A.; English, B.W.; Shokhirev, M.N.; Sinclair, D.A.; Cuellar, T.L. Human age reversal: Fact or fiction? *Ageing Cell* **2022**, *21*, e13664. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
2. Elguezabal-Rodelo, R.; Ochoa-Précoma, R.; Vazquez-Marroquin, G.; Porchia, L.M.; Montes-Arana, I.; Torres-Rasgado, E.; Méndez-Fernández, E.; Pérez-Fuentes, R.; Gonzalez-Mejia, M.E. Metabolic age correlates better than chronological age with waist-to-height ratio, a cardiovascular risk index. *Med. Clin.* **2021**, *157*, 409–417. (In English, Spanish). [[CrossRef](#)] [[PubMed](#)]
3. Oblak, L.; van der Zaag, J.; Higgins-Chen, A.T.; Levine, M.E.; Boks, M.P. A systematic review of biological, social and environmental factors associated with epigenetic clock acceleration. *Ageing Res. Rev.* **2021**, *69*, 101348. [[CrossRef](#)] [[PubMed](#)]
4. Bouchard, C.; Tremblay, A.; Després, J.P.; Nadeau, A.; Lupien, P.J.; Thériault, G.; Dussault, J.; Moorjani, S.; Pinault, S.; Fournier, G. The response to long-term overfeeding in identical twins. *N. Engl. J. Med.* **1990**, *322*, 1477–1482. [[CrossRef](#)] [[PubMed](#)]

5. Ko, S.H.; Jung, Y. Energy Metabolism Changes and Dysregulated Lipid Metabolism in Postmenopausal Women. *Nutrients* **2021**, *13*, 4556. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
6. Speakman, J.R.; Selman, C. Physical activity and resting metabolic rate. *Proc. Nutr. Soc.* **2003**, *62*, 621–634. [[CrossRef](#)] [[PubMed](#)]
7. Mathewson, S.L.; Gordon, A.L.; Smith, K.; Atherton, P.J.; Greig, C.A.; Phillips, B.E. Determining the Influence of Habitual Dietary Protein Intake on Physiological Muscle Parameters in Youth and Older Age. *Nutrients* **2021**, *13*, 3560. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
8. Roberts, S.B.; Dallal, G.E. Energy requirements and aging. *Public Health Nutr.* **2005**, *8*, 1028–1036. [[CrossRef](#)] [[PubMed](#)]
9. Zampino, M.; AlGhatrif, M.; Kuo, P.L.; Simonsick, E.M.; Ferrucci, L. Longitudinal Changes in Resting Metabolic Rates with Aging Are Accelerated by Diseases. *Nutrients* **2020**, *12*, 3061. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
10. Zavros, A.; Andreou, E.; Aphas, G.; Bogdanis, G.C.; Sakkas, G.K.; Roupa, Z.; Giannaki, C.D. The Effects of Zinc and Selenium Co-Supplementation on Resting Metabolic Rate, Thyroid Function, Physical Fitness, and Functional Capacity in Overweight and Obese People under a Hypocaloric Diet: A Randomized, Double-Blind, and Placebo-Controlled Trial. *Nutrients* **2023**, *15*, 3133. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
11. Saito, M.; Matsushita, M.; Yoneshiro, T.; Okamatsu-Ogura, Y. Brown Adipose Tissue, Diet-Induced Thermogenesis, and Thermogenic Food Ingredients: From Mice to Men. *Front. Endocrinol.* **2020**, *11*, 222. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
12. Kruse, M.; Hornemann, S.; Ost, A.C.; Frahn, T.; Hoffmann, D.; Busjahn, A.; Osterhoff, M.A.; Schuppeli, B.; Pfeiffer, A.F.H. An Isocaloric High-Fat Diet Regulates Partially Genetically Determined Fatty Acid and Carbohydrate Uptake and Metabolism in Subcutaneous Adipose Tissue of Lean Adult Twins. *Nutrients* **2023**, *15*, 2338. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
13. Stefanaki, C.; Pervanidou, P.; Boschiero, D.; Chrousos, G.P. Chronic stress and body composition disorders: Implications for health and disease. *Hormones* **2018**, *17*, 33–43. [[CrossRef](#)] [[PubMed](#)]
14. Oster, H. The interplay between stress, circadian clocks, and energy metabolism. *J. Endocrinol.* **2020**, *247*, R13–R25. [[CrossRef](#)] [[PubMed](#)]
15. Deng, H.B.; Tam, T.; Zee, B.C.; Chung, R.Y.; Su, X.; Jin, L.; Chan, T.C.; Chang, L.Y.; Yeoh, E.K.; Lao, X.Q. Short Sleep Duration Increases Metabolic Impact in Healthy Adults: A Population-Based Cohort Study. *Sleep* **2017**, *40*, zsx130. [[CrossRef](#)] [[PubMed](#)]
16. Bonanno, L.; Metro, D.; Papa, M.; Finzi, G.; Maviglia, A.; Sottile, F.; Corallo, F.; Manasseri, L. Assessment of sleep and obesity in adults and children: Observational study. *Medicine* **2019**, *98*, e17642. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
17. GBD 2021 Causes of Death Collaborators. Global burden of 288 causes of death and life expectancy decomposition in 204 countries and territories and 811 subnational locations, 1990–2021: A systematic analysis for the Global Burden of Disease Study 2021. *Lancet* **2024**, *403*, 2100–2132, Erratum in: *Lancet* **2024**, *403*, 1988. [[CrossRef](#)] [[PubMed](#)]
18. Konieczna, J.; Ruiz-Canela, M.; Galmes-Panades, A.M.; Abete, I.; Babio, N.; Fiol, M.; Martín-Sánchez, V.; Estruch, R.; Vidal, J.; Buil-Cosiales, P.; et al. An Energy-Reduced Mediterranean Diet, Physical Activity, and Body Composition: An Interim Subgroup Analysis of the PREDIMED-Plus Randomized Clinical Trial. *JAMA Netw. Open* **2023**, *6*, e2337994. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
19. Saavedra, F.; Fernandes, H.M. Health-related outcomes of strength training in older adults. *Acad. J. Health Sci.* **2022**, *37*, 157–163. [[CrossRef](#)]
20. Celada Roldana, C.; López Díez, J.; Cerezuela, M.A.; Rider, F.; Tárraga Marcos, A.; Tárraga López, P.J.; López González, A.A.; Ramírez Manent, J.I. Cardiovascular effects of a nutritional educational intervention in diabetic patients with poor control. *Acad. J. Health Sci.* **2023**, *38*, 57–65. [[CrossRef](#)]
21. Gómez-Ambrosi, J.; González-Crespo, I.; Catalán, V.; Rodríguez, A.; Moncada, R.; Valentí, V.; Romero, S.; Ramírez, B.; Silva, C.; Gil, M.J.; et al. Clinical usefulness of abdominal bioimpedance (ViScan) in the determination of visceral fat and its application in the diagnosis and management of obesity and its comorbidities. *Clin. Nutr.* **2018**, *37*, 580–589. [[CrossRef](#)] [[PubMed](#)]
22. Gordito Soler, M.; López-González, Á.A.; Vallejos, D.; Martínez-Almoyna Rifá, E.; Vicente-Herrero, M.T.; Ramírez-Manent, J.I. Usefulness of Body Fat and Visceral Fat Determined by Bioimpedanciometry versus Body Mass Index and Waist Circumference in Predicting Elevated Values of Different Risk Scales for Non-Alcoholic Fatty Liver Disease. *Nutrients* **2024**, *16*, 2160. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
23. Bendavid, I.; Lobo, D.N.; Barazzoni, R.; Cederholm, T.; Coëffier, M.; de van der Schueren, M.; Fontaine, E.; Hiesmayr, M.; Laviano, A.; Pichard, C.; et al. The centenary of the Harris-Benedict equations: How to assess energy requirements best? Recommendations from the ESPEN expert group. *Clin. Nutr.* **2021**, *40*, 690–701. [[CrossRef](#)] [[PubMed](#)]
24. Di Vincenzo, J.D.; O'Brien, L.; Jacobs, I.; Jawad, M.Y.; Ceban, F.; Meshkat, S.; Gill, H.; Tabassum, A.; Phan, L.; Badulescu, S.; et al. Indirect Calorimetry to Measure Metabolic Rate and Energy Expenditure in Psychiatric Populations: A Systematic Review. *Nutrients* **2023**, *15*, 1686. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
25. Galgani, J.E.; Fernández-Verdejo, R. Pathophysiological role of metabolic flexibility on metabolic health. *Obes. Rev.* **2021**, *22*, e13131. [[CrossRef](#)] [[PubMed](#)]
26. la Torre, A.; Lo Vecchio, F.; Greco, A. Epigenetic Mechanisms of Aging and Aging-Associated Diseases. *Cells* **2023**, *12*, 1163. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
27. Żelaźniewicz, A.; Nowak, J.; Łacka, P.; Pawłowski, B. Facial appearance and metabolic health biomarkers in women. *Sci. Rep.* **2020**, *10*, 13067. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
28. Stewart, A.; Marfell-Jones, M.; Olds, T.; Ridder, H. *International Standards for Anthropometric Assessment*; International Society for the Advancement of Kinanthropometry–ISAK: Lower Hutt, New Zealand, 2011.

29. Fang, H.; Berg, E.; Cheng, X.; Shen, W. How to best assess abdominal obesity. *Curr. Opin. Clin. Nutr. Metab. Care* **2018**, *21*, 360–365. [[CrossRef](#)]
30. Riutord-Sbert, P.; Riutord-Fe, B.; Riutord-Fe, N.; Arroyo-Bote, S.; López González, A.A.; Ramírez-Manent, J.I. Influence of physical activity and mediterranean diet on the values of different scales of overweight and obesity. *Acad. J. Health Sci.* **2022**, *37*, 21–28.
31. Martínez Jover, A.; López-González, A.A.; Tomás Gil, P.; Coll Villalonga, J.L.; Martí Lliteras, P.; Ramírez-Manent, J.I. Variables influencing the appearance of metabolic syndrome with three different definitions in 418,343 spanish workers. *Acad. J. Health Sci.* **2023**, *38*, 129–135. [[CrossRef](#)]
32. Craig, C.L.; Marshall, A.L.; Sjöström, M.; Bauman, A.E.; Booth, M.L.; Ainsworth, B.E.; Pratt, M.; Ekelund, U.L.F.; Yngve, A.; Sallis, J.F.; et al. International physical activity questionnaire: 12-country reliability and validity. *Med. Sci. Sports Exerc.* **2003**, *35*, 1381–1395. [[CrossRef](#)] [[PubMed](#)]
33. Domingo-Salvany, A.; Bacigalupe, A.; Carrasco, J.M.; Espelt, A.; Ferrando, J.; Borrell, C.; del Grupo de Determinantes Sociales de Sociedad Española de Epidemiología. Propuestas de clase social neoweberiana y neomarxista a partir de la Clasificación Nacional de Ocupaciones 2011 [Proposals for social class classification based on the Spanish National Classification of Occupations 2011 using neo-Weberian and neo-Marxist approaches]. *Gac. Sanit.* **2013**, *27*, 263–272. (In Spanish) [[CrossRef](#)] [[PubMed](#)]
34. Ramírez Gallegos, I.; Marina Arroyo, M.; López-González, Á.A.; Vicente-Herrero, M.T.; Vallejos, D.; Sastre-Alzamora, T.; Ramírez-Manent, J.I. The Effect of a Program to Improve Adherence to the Mediterranean Diet on Cardiometabolic Parameters in 7034 Spanish Workers. *Nutrients* **2024**, *16*, 1082. [[CrossRef](#)] [[PubMed](#)]
35. Elliott, M.L.; Caspi, A.; Houts, R.M.; Ambler, A.; Broadbent, J.M.; Hancox, R.J.; Harrington, H.; Hogan, S.; Keenan, R.; Knodt, A.; et al. Disparities in the pace of biological aging among midlife adults of the same chronological age have implications for future frailty risk and policy. *Nat. Aging* **2021**, *1*, 295–308. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
36. Flynn, M.G.; Markofski, M.M.; Carrillo, A.E. Elevated Inflammatory Status and Increased Risk of Chronic Disease in Chronological Aging: Inflamm-aging or Inflamm-inactivity? *Aging Dis.* **2019**, *10*, 147–156. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
37. Rattan, S.I. Aging is not a disease: Implications for intervention. *Aging Dis.* **2014**, *5*, 196–202. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
38. Wensink, M.J.; van Heemst, D.; Rosing, M.P.; Westendorp, R.G. The maintenance gap: A new theoretical perspective on the evolution of aging. *Biogerontology* **2012**, *13*, 197–201. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
39. Mauvais-Jarvis, F.; Bairey Merz, N.; Barnes, P.J.; Brinton, R.D.; Carrero, J.J.; DeMeo, D.L.; De Vries, G.J.; Epperson, C.N.; Govindan, R.; Klein, S.L.; et al. Sex and gender: Modifiers of health, disease, and medicine. *Lancet* **2020**, *396*, 565–582, Erratum in: *Lancet* **2020**, *396*, 668. [[CrossRef](#)] [[PubMed](#)]
40. Marlatt, K.L.; Pitynski-Miller, D.R.; Gavin, K.M.; Moreau, K.L.; Melanson, E.L.; Santoro, N.; Kohrt, W.M. Body composition and cardiometabolic health across the menopause transition. *Obesity* **2022**, *30*, 14–27. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
41. Kanellakis, S.; Skoufas, E.; Simitsopoulou, E.; Migdanis, A.; Migdanis, I.; Prelorentzou, T.; Louka, A.; Moschonis, G.; Bountouvi, E.; Androutsos, O. Changes in body weight and body composition during the menstrual cycle. *Am. J. Hum. Biol.* **2023**, *35*, e23951. [[CrossRef](#)] [[PubMed](#)]
42. Stringhini, S.; Carmeli, C.; Jokela, M.; Avendaño, M.; Muennig, P.; Guida, F.; Ricceri, F.; d’Errico, A.; Barros, H.; Bochud, M.; et al. Socioeconomic status and the 25 × 25 risk factors as determinants of premature mortality: A multicohort study and meta-analysis of 1.7 million men and women. *Lancet* **2017**, *389*, 1229–1237, Erratum in: *Lancet* **2017**, *389*, 1194. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
43. Woodward, E.N.; Walsh, J.L.; Senn, T.E.; Carey, M.P. Positive social interaction offsets impact of low socioeconomic status on stress. *J. Natl. Med. Assoc.* **2018**, *110*, 371–377. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
44. Marmot, M.; Friel, S.; Bell, R.; Houweling, T.A.; Taylor, S.; Commission on Social Determinants of Health. Closing the gap in a generation: Health equity through action on the social determinants of health. *Lancet* **2008**, *372*, 1661–1669. [[CrossRef](#)] [[PubMed](#)]
45. Montero Muñoz, N.; López-González, A.A.; Tomás-Gil, P.; Martínez Jover, A.; Paublini, H.; Ramírez-Manent, J.I. Relationship between sociodemographic variables and tobacco consumption with vascular age values using the Framingham model in 336,450 spanish workers. *Acad. J. Health Sci.* **2023**, *38*, 61–66. [[CrossRef](#)]
46. Behl, T.A.; Stamford, B.A.; Moffatt, R.J. The Effects of Smoking on the Diagnostic Characteristics of Metabolic Syndrome: A Review. *Am. J. Lifestyle Med.* **2022**, *17*, 397–412. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
47. Park, S.; Kim, S.G.; Lee, S.; Kim, Y.; Cho, S.; Kim, K.; Kim, Y.C.; Han, S.S.; Lee, H.; Lee, J.P.; et al. Causal linkage of tobacco smoking with ageing: Mendelian randomization analysis towards telomere attrition and sarcopenia. *J. Cachexia Sarcopenia Muscle* **2023**, *14*, 955–963. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
48. Larsson, S.C.; Burgess, S. Appraising the causal role of smoking in multiple diseases: A systematic review and meta-analysis of Mendelian randomization studies. *EBioMedicine* **2022**, *82*, 104154. [[CrossRef](#)] [[PubMed](#)]
49. Kelly, R.S.; Kelly, M.P.; Kelly, P. Metabolomics, physical activity, exercise and health: A review of the current evidence. *Biochim. Biophys. Acta Mol. Basis Dis.* **2020**, *1866*, 165936. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
50. Terada, T.; Boulé, N.G.; Forhan, M.; Prado, C.M.; Kenny, G.P.; Prud’homme, D.; Ito, E.; Sigal, R.J. Cardiometabolic risk factors in type 2 diabetes with high fat and low muscle mass: At baseline and in response to exercise. *Obesity* **2017**, *25*, 881–891. [[CrossRef](#)] [[PubMed](#)]
51. Burini, R.C.; Anderson, E.; Durstine, J.L.; Carson, J.A. Inflammation, physical activity, and chronic disease: An evolutionary perspective. *Sports Med. Health Sci.* **2020**, *2*, 1–6. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]

52. Krebs-Smith, S.M.; Pannucci, T.E.; Subar, A.F.; Kirkpatrick, S.I.; Lerman, J.L.; Tooze, J.A.; Wilson, M.M.; Reedy, J. Update of the Healthy Eating Index: HEI-2015. *J. Acad. Nutr. Diet.* **2018**, *118*, 1591–1602, Erratum in: *J. Acad. Nutr. Diet.* **2019**, *119*, 1759. [[CrossRef](#)] [[PubMed](#)]
53. Dayi, T.; Ozgoren, M. Effects of the Mediterranean diet on the components of metabolic syndrome. *J. Prev. Med. Hyg.* **2022**, *63* (Suppl. S3), E56–E64. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
54. Guasch-Ferré, M.; Willett, W.C. The Mediterranean diet and health: A comprehensive overview. *J. Intern. Med.* **2021**, *290*, 549–566. [[CrossRef](#)] [[PubMed](#)]
55. Obeid, C.A.; Gubbels, J.S.; Jaalouk, D.; Kremers, S.P.J.; Oenema, A. Adherence to the Mediterranean diet among adults in Mediterranean countries: A systematic literature review. *Eur. J. Nutr.* **2022**, *61*, 3327–3344. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
56. Herrera-Ramos, E.; Tomaino, L.; Sánchez-Villegas, A.; Ribas-Barba, L.; Gómez, S.F.; Wärnberg, J.; Osés, M.; González-Gross, M.; Gusi, N.; Aznar, S.; et al. Trends in Adherence to the Mediterranean Diet in Spanish Children and Adolescents across Two Decades. *Nutrients* **2023**, *15*, 2348. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
57. Nations, U. World Population Prospects: The 2017 Revision, Key Findings and Advance Tables. Working Paper No ESA/P/WP/248. 2017. Available online: https://population.un.org/Wpp/Publications/Files/Wpp2017_Keyfindings.Pdf (accessed on 12 October 2024).
58. Li, C.W.; Yu, K.; Shyh-Chang, N.; Jiang, Z.; Liu, T.; Ma, S.; Luo, L.; Guang, L.; Liang, K.; Ma, W.; et al. Pathogenesis of sarcopenia and the relationship with fat mass: Descriptive review. *J. Cachexia Sarcopenia Muscle* **2022**, *13*, 781–794. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
59. Blüher, M. Obesity: Global epidemiology and pathogenesis. *Nat. Rev. Endocrinol.* **2019**, *15*, 288–298. [[CrossRef](#)] [[PubMed](#)]
60. Brunjes, D.L.; Kennel, P.J.; Christian Schulze, P. Exercise capacity, physical activity, and morbidity. *Heart Fail. Rev.* **2017**, *22*, 133–139. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]
61. de Meijer, C.; Koopmanschap, M.; D’Uva, T.B.; van Doorslaer, E. Determinants of long-term care spending: Age, time to death or disability? *J. Health Econ.* **2011**, *30*, 425–438. [[CrossRef](#)] [[PubMed](#)]
62. WHO. *Saving Lives, Spending Less: The Case for Investing in Noncommunicable Diseases*; World Health Organization: Geneva, Switzerland, 2021. Available online: <https://iris.who.int/bitstream/handle/10665/350449/9789240041059-eng.pdf?sequence=1> (accessed on 12 October 2024).
63. Ding, D.; Lawson, K.D.; Kolbe-Alexander, T.L.; Finkelstein, E.A.; Katzmarzyk, P.T.; van Mechelen, W.; Pratt, M.; Lancet Physical Activity Series 2 Executive Committee. The economic burden of physical inactivity: A global analysis of major non-communicable diseases. *Lancet* **2016**, *388*, 1311–1324. [[CrossRef](#)] [[PubMed](#)]
64. *Purchasing Power Parities—Frequently Asked Questions*; OECD: Paris, France, 2021.
65. Santos, A.C.; Willumsen, J.; Meheus, F.; Ilbawi, A.; Bull, F.C. The cost of inaction on physical inactivity to public health-care systems: A population-attributable fraction analysis. *Lancet Glob. Health* **2023**, *11*, e32–e39. [[CrossRef](#)] [[PubMed](#)] [[PubMed Central](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.